

Estimated Global Mortality from Present to 2100 from Climate Change

William N. Rom M.D., MPH

Sol and Judith Bergstein Professor of Medicine and Environmental Medicine
Division of Pulmonary, Critical Care, and Sleep Medicine
Departments of Medicine and Environmental Medicine
New York University School of Medicine
NYU College of Global Public Health
550 1st Ave, New York, NY 10016, USA.

Abstract

Introduction. Climate change mortality emanates from increased heat, air pollution, extreme weather/damage to coastal cities, coral death and loss of fisheries, and food insecurity/population growth and migration.

Methods. The purpose of this paper is to propose estimates of mortality 2018-2100 with the objective to increase observations in order to develop methodology to accurately predict climate change mortality.

Results. The past three years have broken records for global warming due to continued release of greenhouse gases. CO₂ is the primary driver, and is a long-lived greenhouse gas with one-third remaining in the atmosphere after a century of its original release. The Intergovernmental Panel on Climate Change (IPCC) Representative Concentration Pathway 2.6 estimated mortality is 13.5 million versus 106.5 million for RCP 8.5.

Conclusion. Business-as-usual with continued release of 40 billion tons of CO₂ may increase global mean temperature by 4.8±4 °C increasing mortality to 2 million deaths/year for the period 2050-2100 from the WHO estimate of 250,000/year 2030-2050.

Introduction.

Global warming became a term to describe the global greenhouse effect following Dr. James Hansen's testimony June 23, 1988, before the U.S. Senate; he raised such alarm that "Climate Change" was subsequently used which also broadened the scope of ecological concern (Hansen 2013). The concepts of a greenhouse effect from anthropogenic CO₂ release were recognized by Svante Arrhenius in 1896. Measurements of CO₂ and global temperature from many sources led to substantiation of the theory, and by the end of the 1990s, climate change gained traction because every year reached record highs for mean global temperature (Rom 2014). The last three years 2014, 2015, and 2016 broke the global temperature record each year (Karl 2015). With an increase of global mean temperature of 1 °C from the first 500-600 gigatons of CO₂ released, there is a carbon budget of 150-1050 Gt remaining to keep from exceeding the 1.5° C target global temperature increase agreed to by the 195 nations signing the Paris Climate Agreement in 2015. The World Health Organization has estimated global mortality from climate change to be 150,000 deaths/year increasing to 250,000 deaths/year by 2030 (Hales 2014). The purpose of this Commentary is to estimate global mortality due to climate change from the present to 2100 using the Intergovernmental Panel on Climate Change (IPCC) Representative Concentration Pathway (RCP) of 2.6 Watts/cm² versus Business-as Usual RCP of 8.5 Watts/cm² (Brown 2014, Hansen 2016). There currently is no methodology for accurately projecting mortality, and this Commentary hopefully will encourage development of biostatistical and computational methods. Currently estimates vary widely, e.g. Hurricane Maria officially caused 64 deaths in Puerto Rico although news reports estimate 1,052 deaths.

Moreover, dengue, spread by *Aedes aegypti* mosquitoes whose range is expected to increase dramatically 2050-2100 due to a warmer and wetter mid-latitudes, has been estimated to cause 12,500 deaths annually (Liu-Helmersson 2014). In forecasting the future, attribution of health effects due to climate change will increase as temperature and humidity also increase.

The Physical Science of Global Warming.

The amount of CO₂ in Earth's atmosphere has been chronicled by the Keeling curve representing data collected from the summit of Mauna Loa in Hawaii since 1959, the International Geophysical Year (Rom 2014). In 1959 the level was approximately 310 ppm; this level was higher compared to the 250 ppm measured in bubbles in ice cores from Greenland or Antarctica, and going back 400,000 to 800,000 years (Luthi 2008). From 1959 to 2016 the atmospheric CO₂ has increased from 310 to 400 ppm (~43%). Importantly, the annual rise or rate of increase has jumped from less than 2 ppm to more than 3 ppm (over the past two years) (Zeebe 2016). CO₂ forms a greenhouse gas layer around the earth that allows sunlight to pass through, and then reflects infrared radiation back to the surface of the earth causing surface heating. Methane, CH₄, is approximately 22 times more potent as a greenhouse gas than CO₂, and may be 80 times more potent over a 20-year time frame (Montzka 2011). CH₄ in parts per billion is emitted from landfills, fracking, conventional natural gas, and oil wells, leaks in pipelines, wetlands, emissions from livestock, and melting permafrost. Additionally, there are other strong greenhouse gases, e.g. nitrous oxide emitted from rice fields and fertilizers. Carbon black is emitted by diesel engines and biomass cooking in developing countries. Several gases used as refrigerants and in air conditioners such as hydrofluorocarbons, have a global warming potential up to 2000 times that of CO₂ (Molina 2009). The source of carbon pollution is anthropogenic: oil used in industry and transportation, natural gas and coal used in electricity generation and steel manufacture, cement production, and buildings in cities (IPCC 2014). The increase in the anthropogenic greenhouse gas releases primarily occurred over the past 50 years, and parallels the increase in CO₂ measurements, i.e. the Earth is out of energy balance (Karl 2009, USGCRP 2014). Importantly, CO₂ pollution is different from other air pollutants while particulate matter or ozone varies by hour or day or season, CO₂ is persistent and cumulative (Driscoll 2015). After 100 years, 33% remains in the atmosphere, and after a millennia, 20% remains (Solomon 2009). CO₂ is part of the carbon cycle with major sinks in the oceans and soil. The world's forests remove CO₂ from the atmosphere in the process of photosynthesis and the Earth's energy balance is further upset by deforestation (Levinson 2014).

The amount of carbon dioxide emissions we can emit while still having a likely chance of limiting global temperature to 2° C above pre-industrial levels after 565 Gigatons (Gt) already released is 150-790 Gt more (McKibben 2012). At global annual emissions of ~40 Gt CO₂ we will exhaust the carbon budget in 3-22 years for stabilization at 2° C. In the IPCC Representative Concentration Pathway (RCP) 8.5 (Business-as-Usual) CO₂ emissions approximately double by 2050 and increase by three-fold by 2100 with corresponding increases in CO₂ beyond 900 ppm and global temperature increase of as much as 4.8±4° C (Brown 2017). Unknown in this scenario would be release of CO₂ and methane from permafrost thawing across the tundra regions of Canada, Russia, and Alaska, and de-stabilization of Antarctic and Greenland glaciers.

The purpose of this paper is to encourage methodologies to accurately forecast future mortality from various scenarios of global heating and the five topics contributing to mortality outlined below. The temperatures at the end of this period (2080-2100) will be outside the range most living species have ever experienced (Moritz 2013). Mortality under RCP 8.5 is predicted to increase after 2050 to 2 million deaths/year (an 8-fold increase) and persist at least at that level through 2100 (cumulative 106.5 million deaths) due to global heating compared to 13.5 million (250,000 deaths/year) under RCP 2.6 (Watts 2015, 2017). The five topical areas that will feature climate-related mortality are discussed below (Watts 2015, Watts 2017, Whitmee 2015).

Heat waves may contribute approximately 10% to the 2050-2100 RCP 8.5 mortality (Gasparini 2015). Heat waves will increase in frequency from once per 500 years to annual events with wide regional differences. In 2003 there was a heat wave in France where temperatures exceeded the norm by 11-12° C for 9 consecutive days. There were 15,000 deaths in France; 32,000 in western Europe, and 70,000 across Europe during this period (Fouillet 2006). These increases illustrate how climate change mortality will impact large geographic areas. Mortality was age-related and was 45% higher in women than in men >45 years. Mortality was increased in widowed, single and divorced people, and lack of air conditioning was a major cause.

A significant increase in deaths was related to heatstroke, hyperthermia, and dehydration; heart failure, chronic respiratory diseases and stroke were markedly increased, and psychiatric disorders, especially depression increased (Hales 2014). More recently heat waves have been recorded in Moscow (associated with wild fires) and New Delhi (Mazdiyasn 2017). Using a probabilistic model, an increase of mean temperatures over India of 0.5 °C corresponded to a 146% increase in the probability of heat-related mortality events of more than 100 people (Mazdiyasn 2017). There are heat islands surrounding urban areas from the black asphalt pavement, buildings, and lack of green space (Greene 2011). During heat waves, there is a lack of cooling at night since the heat does not dissipate. Heat waves are accompanied by dry periods leading to a lack of moisture in neighboring forests and bushlands. These can easily catch fire from humans or lightning strikes causing extensive smoke. In the United States, the western fire season has been extended 78 days since 1970, and the average duration of fires has increased five-fold. Epidemiological studies have shown that an increase of 10 µg/m³ in PM₁₀ from wildfires results in approximately 1% increase in non-accidental mortality (Whitmee 2015). During Australian brushfires overall mortality rose 5%, and hospital admissions for respiratory illnesses increased from 3-5% (Johnston 2011). New South Wales in Australia had a brutal heat wave in the summer of 2017 with temperatures reaching 113° F; World Weather Attribution found these maximum temperatures were 10 times more likely compared to a century ago before anthropogenic global heating (Whitmee 2015). In addition to heat waves, global heating will increase mortality for certain diseases including arthropod-borne diseases as the mosquito range increases for dengue, malaria, zika, yellow fever, and chikungunya; water-borne illnesses due to vibrio, cryptosporidia; and rodent-borne diseases including lyme disease (Rom 2014). In addition, heat has an adverse effect on pregnancy outcomes (Watts 2017). More recently, heat related mortality occurs apart from heat waves due to lack of adaptation to increasing temperatures (O'Neill 2005, Watts 2017). An estimated 125 million additional vulnerable adults were exposed to heat waves between 2000 and 2016 (Watts 2017).

Mora and colleagues described in “Global risk of deadly heat,” that one-third of the world’s population is currently exposed to a deadly combination of surface temperature exceeding 95 °F and relative humidity for at least 20 days a year (Mora 2017). This will increase in 2100 to 74%, and even under RCP 2.6 will reach 48%. New York City could reach 50 days in the summer, but cities like Jakarta where their consistently warm temperatures would be near the deadly threshold year-around (Coffel 2017). Both ageing of the population and increasing urbanization could aggravate vulnerability to heat. Coffel and colleagues estimated that half of the world’s population would be exposed to wet bulb temperatures that approach and exceed limits of human tolerability; under RCP 8.5 exposure to wet bulb temperatures above 35° C could exceed a million person-days per year by 2080 (Coffel 2017).

2). Air pollution may result in 20% of increased mortality although 7 million deaths/year are attributed from air pollution now (Doherty 2009, Balbus 2016). Currently the Global Burden of Disease estimates 3.5 million deaths/year from outdoor air pollution primarily due to cardiac and respiratory conditions and lung cancer (Hoek 2013, Lelieveld 2015). Another 3.5 million deaths/year occur from indoor air pollution due to biomass cooking (charcoal, dung, coal, wood) affecting primarily women (chronic obstructive pulmonary disease and lung cancer) and children (respiratory bronchiolitis and pneumonia) (Rice 2014). There is overlap between climate change health effects and air pollution health effects with co-benefits accruing from decreases in coal combustion and increased market penetration of renewable energy and electric vehicles (Berman 2012, West 2013, Garcia-Menendez 2015). Black carbon from diesel engines and biomass cooking is a short-lived global heating contributor. The Global Alliance for Clean Cookstoves has a target of 100 million clean cookstoves by 2020. Air pollution should decline 2050-2100 due to phase-out of coal combustion, introduction of electric vehicles, and clean cookstoves (Cromar 2016).

3). Droughts and heat effects on agriculture may cause 40% of the RCP 8.5 mortality 2050-2100 (Ebi 2016, Patz 2014, Egan 2016). The increased heat will cause severe droughts in certain areas, e.g. United States Southwest and California, whereas other areas will experience an increase in precipitation. Temperature has a variable effect on different crop species but higher temperatures will cause yield declines between 2.5% and 10% across a number of agronomic species; extreme high temperatures during the reproductive stage will affect pollen viability, fertilization, and grain or fruit formation (Hatfield 2015). Wheat, corn, and cotton yields have statistical declines between 63% to 70% for high CO₂ emission scenarios (before accounting for positive effects of rising atmospheric CO₂ on crop growth). The global population of 7.3 billion at present will grow to 9.3 billion by 2050 and 11.2 billion by 2100 with a corresponding demand for more food (McMichael 2013).

Higher minimum temperatures reduce grain yield of rice, and extreme maximum temperatures could also reduce rice yield (Hatfield 2015). Heat stress of 4 °C above optimum could lead to crop failure. Under global heating, perennial crops may not have enough hours of cooling to produce fruits, nuts, grapes, etc. requiring innovative strategies for genetic selection and tree and vine growth beyond 2050. Temperature effects interact with soil water status to increase the negative effects on grain production. The greatest risk for drought and food scarcity will be in Africa's Sahel, Ethiopia and Sudan, the Middle East, south and east Asia especially India, China, Russia and Central and South America (Park 2018). The Asian Development Bank estimates a 4.6°C rise in global temperature in 2100 with an 8.8% loss in annual domestic gross domestic product in 6 countries of South Asia including India.

4). Loss of coastal coral reefs due to global heating may result in a surprising 10% increase in mortality in coastal communities due to dependence on fish and marine life for sustenance that will dramatically decline (Anthony 2008). Global heating will also result in reductions of other fisheries such as cod off the Maine coast (Pershing 2015). Although 0.1% of the oceans are populated by coral reefs, these ecosystems are responsible for over one-third of fish reproduction. For example, 357 million people live in the Philippines and Indonesia, and 40% of them live in coastal communities that are dependent on the ocean for means of sustenance. Their coral reefs are all under heat stress with only 1% of Philippine reefs in pristine condition. The median return time between pairs of severe bleaching events has diminished steadily since 1980 globally, and is now only 6 years (Hughes 2018). Shortening the coral bleaching recovery time increases likelihood of coral death. 39% of the 40 million Filipinos living on the coasts live in poverty and 36% are reported to be food insecure. The Coral Triangle (Indonesia, Malaysia, East Timor, Philippines, Solomon Islands) encompasses 130 million people and their food, income, and livelihood depend on the coral reefs and fisheries (McIver 2016). They provide 11.3% of global fisheries' productivity and the Melanesian fisheries are worth \$7 billion USD. The Food and Agriculture Organization of the United Nations estimates that climate-related natural disasters doubled over the period 2003-2013 compared to the 1980s causing an estimated \$1.5 trillion USD in economic damages and leaving many vulnerable populations in developing countries exposed to >50% food insecurity.

5). Storms, typhoons, hurricanes, flooding and weather-related events may result in the remaining 20% mortality (Grinsted 2013, Fisher 2015, Stott 2016). Hurricane Sandy in 2012 caused \$60 billion USD in losses, 117 deaths (40 from drowning), and flooded much of lower Manhattan in New York City with a 14-foot storm surge (MMWR 2012, Rom 2013). Hurricanes may increase in intensity, and by 2050 will have increased in frequency, such that a Hurricane Katrina-like event could happen every two years. Hurricane Katrina caused over 1,800 deaths and over \$100 billion USD in damage. Typhoon Haiyan was the deadliest Philippine typhoon on record, killing 6,340 people with 1,061 missing. Haiyan was also the strongest storm recorded at landfall in November 2013 with 1-minute sustained winds reaching 315 km/h (195 mph). Over 11 million people were left homeless in the Philippines (National Academy of Sciences, Engineering and Medicine 2016). Global heating will increase evaporation and cloud formation leading to heavy rainfall and rapid flooding in certain regions of the world (Taylor 2017). Mortality is difficult to estimate from hurricanes, e.g. Hurricane Maria's estimates ranging from 64 to 1,052 in Puerto Rico are still being investigated months later. Coastal flooding will increase due to ocean rise, and high tides will exacerbate coastal damage to homes, infrastructure and cities. Morabia and Benjamin described a new public health priority of rebuilding after climate change-related natural disasters, e.g. from three >\$1 billion USD natural disasters in 1980 to seventeen in 2017 totaling \$306 billion USD (Morabia 2018). The global mean sea level rise has increased from 2.2 mm/year in 1993 to 3.3 mm/year in 2014 as measured by satellite altimetry (Chen 2017). Sea level rise may exceed 2 meters in the RCP 8.5 scenario (DeConto 2016, Koppo 2014). Importantly, this is an issue of environmental justice since the mortality will fall disproportionately upon the poor, elderly, children, women, in low-income countries across Africa, South and East Asia, and South and Central America. Efforts at coastal and housing adaptation can mitigate storm-related mortality in the future as well.

Policy Implications of Climate Change Mortality.

Achieving an 80% reduction in CO₂ emissions by 2050 (RCP 2.6) would require three strategies:

1). *Transitioning to a low-carbon energy system.* This requires an increase in energy efficiency of buildings, appliances, more light-emitting diode (LEDs) in lighting, and conversion to electric vehicles. It is essential to decarbonize electricity by moving to ~55% wind and solar, 17% nuclear, 20% natural gas with carbon capture and storage, and 8% hydroelectric (The White House 2016). Electricity storage in large lithium battery facilities or other technologies, a demand response of the grid, and direct current high voltage transmission for renewable energy will be necessary.

2). *Sequestering carbon through forests and soils and CO₂ removal technologies.* There would be a need for at least a 50 million acres of forest expansion (U.S. alone and more elsewhere), and research to increase soil mass and depth of commodity crops (The White House 2016).

3). *There is a need for a policy to reduce non-CO₂ emissions which contribute ~20% of global heating, e.g. methane, nitrous oxide, and hydrofluorocarbons* (Pacala 2004).

An example was published by MacDonald and colleagues using a model of U.S. electricity generating capacity to achieve a 78% reduction of CO₂ emissions relative to 1990 and building renewable energy primarily wind and solar PV plants (MacDonald 2016). It included 523 gigawatts (GW) of wind (22 MW offshore), 371 GW of solar PV, 461 GW of nuclear, and 74 GW of hydroelectric with the remaining natural gas for a total of 1,529 GW of installed capacity. Their model focused on weather where wind blows on the U.S. Great Plains and the sun shines in the Southwest, with High Voltage Direct Current transmission lines transporting power to markets via 32 nodes. Their model predicted a growth of wind by a factor of eight and solar PV by a factor of 62. Although they did not discuss offshore wind, wind turbines there could be larger (6 MW) reducing the number needed by half, and can offer 33% more power due to more frequent and stronger wind over the sea. In addition, offshore wind is close to major coastal cities, and transport is less likely over private land (Jacobson 2016).

In summary, global heating is upon us, and an estimate of 106.5 million people may perish globally between 2018 and 2100 under the RCP 8.5 scenario compared to 13.5 million with the RCP 2.6 scenario (Markandya 2009). The period 2018-2050 is critical to advance towards a global clean energy revolution (Patz 2014, Rom 2008, Horton 2009, Millar 2017, Watts 2015, 2017).

References.

1. Anthony KRN, Kline DI, Diaz-Palido, Dove S, Hoegh-Guldberg O. 2008. Ocean acidification causes bleaching and productivity loss in coral reef builders. *Proc Natl Acad Sci USA*. 105: 17442-6.
2. Balbus JM, Tart KGT, Dilworth CH, Birnbaum LS. 2016. Changing the climate of respiratory clinical practice. Insights from the 2016 Climate and Health Assessment of the U.S. Global Change Research Program. *Am J Respir Crit Care Med* 13: 1202-1204.
3. Berman JD, Fann N, Hollingsworth JW, Pinkerton KE, Rom WN, Szema AM, Breyse PN, White RH, Curriero FC. 2012. Health benefits from large-scale ozone reduction in the United States. *Environmental Health Perspectives* 2012; 120: 1404-1410. doi: 10.1289/ehp.1104851.
4. Brown H and Spickett J. 2014. Health consequence scales for use in health impact assessments of climate change. *Int J Environ Res Public Health* 11: 9607-9620.
5. Brown PT and Caldeira K. 2017. Greater future global warming inferred from Earth's recent energy budget. *Nature* 552: 45-50. doi:10.1038/nature24672
6. Chen X, Zhang X, Church JA, Watson CS, King MA, Monselesan D, Legresy B, Harig C. 2017. The increasing rate of global mean sea-level rise during 1993-2014. *Nature Climate Change* 7: 492-495. DOI: 10.1038/NCLIMATE3325.
7. Coffel ED, Horton RM, Sherbinin A. 2017. Temperature and humidity projections of a rapid rise in global heat stress exposure during the 21st century. *Environmental Research Letters*. 13: 014001. Doi.org/10.1088/1748-9326/aaa00e
8. Cromar KR, Gladson LA, Perlmutter LD, Ghazipura M, Ewart GW. 2016. Estimated excess morbidity and mortality caused by air pollution above American Thoracic Society-recommended standards, 2011-2013. *Ann Am Thoracic Soc* 13(8): 1195-1201.
9. DeConto RM and Pollard D. 2016. Contribution of Antarctica to past and future sea-level rise. *Nature* 531: 591-597. doi:10.1038/nature17145

10. Doherty RM, Heal MR, Wilkinson P, Pattenden S, Vieno M, Armstrong B, Atkinson R, Chalabi Z, Kovats S, Milojevic A, Stevenson DS. 2009. Current and future climate- and air pollution-mediated impacts on human health. *Environmental Health* 8(Suppl 1): S8.
11. Driscoll CT, Buonocore JJ, Levy JI, Lambert KF, Burtraw D, Reid SB, Fakhraei H, Schwartz J. 2015. U.S. power plant carbon standards and clean air and health co-benefits. *Nature Climate Change* 5: 535–540. doi:10.1038/NCLIMATE2598.
12. Ebi K and Bowen KJ. 2016. Extreme events as sources of health vulnerability: Drought as an example. *Weather and Climate Extremes*. 11: 95–102. doi:10.1016/j.wace.2015.10.001.
13. Egan PJ and Mullin M. 2016. Recent improvement and projected worsening of weather in the United States. *Nature* 532: 357-360. doi:10.1038/nature17441.
14. Fischer EM and Knutti R. 2015. Anthropogenic contribution to global occurrence of heavy-precipitation and high-temperature extremes. *Nature Climate Change* 5: 560-564. Doi: org/10.1038/nclimate2617.
15. Fouillet A, Rey G, Laurent F, Pavillon G, Bellec S, Guihenneuc-Jouyau C, Clavel J, Jouglu E, Hemon D. 2006. Excess mortality related to the August 2003 heat wave in France. *Int Arch Occup Environ Health* 80: 16-24.
16. Garcia-Menendez F, Saari RK, Monier E, Selin NE. 2015. U.S. air quality and health benefits from avoided climate change under greenhouse gas mitigation. *Environ Sci Technol* 49: 7580-7588.
17. Gasparini A, Guo Y, Hashizume M, Kinney PL, Petkova EP, Lavigne E, Zanobetti A, Schwartz JD, Tobias A, Leone M, Tong S, Honda Y, Kim H, Armstrong BG. 2015. Temporal variation in heat-mortality associations: A multicountry study. *Environ Health Persp* 123: 1200-1207.
18. Greene S, Kalkstein LS, Mills DM, Samenow J. 2011. An examination of climate change on extreme heat events and climate-change mortality relationships in large U.S. cities. *Weather Clim Soc* 3: 281-291.
19. Grinsted A, Moore JC, and Jevrejeva S. 2013. Projected Atlantic hurricane surge threat from rising temperatures. *Proc Natl Acad Sci USA*. 110: 5369-5373. doi 10.1073/pnas.1209980110.
20. Hales S, Kovats S, Lloyd S, Campbell-Lendrum D. 2014. Quantitative risk assessment of the effects of climate change on selected causes of death 2030s and 2050s. Geneva: World Health Organization, ISBN 978 92 4 150769 1
21. Hansen J, Kharecha P, Sato M, Masson-Delmotte V, Ackerman F, Beerling DJ. 2013. Assessing “dangerous climate change”: Required reduction of carbon emissions to protect young people, future generations and nature. *PLoS ONE* 8(12), e81648. Doi:10.1371/journal.pone.0081648
22. Hansen J, Sato M, Hearty P, Ruedy R, Kelley M, Masson-Delmotte V, Russell G, Tseliodis G, Cao J, Rignot E, Velicogna I, Tormey B, Donovan B, Kandiano E, Schuckman K, Kharecha P, Legrande AN, Bauer M, Lo K-W. 2016. Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2° C global warming could be dangerous. *Atmos Chem Phys* 16: 3761-3812. Doi: 10.5194/acp-16-3761-2016
23. Hatfield JL and Prueger JH. 2015. Temperature extremes: Effect on plant growth and development. *Weather and Climate Extremes* 104: 4-10. Doi: 10.1016/j.wace.2015.08.001
24. Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, Kaufman JD. Long-term air pollution exposure and cardio-respiratory mortality: A review. *Environmental Health* 2013; 12: 43-58.
25. Horton R. Health and Climate Change. The climate dividend. *The Lancet* 2009; DOI:10.1016/S0140-6736(09)61994-2.
26. Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC, Claar DC, Eakin CM, Gilmour JF, Graham NAJ, Harrison H, Hobbs J-PA, Hoey AS, Hoogenboom M, Lowe RJ, McCulloch MT, Pandolfi JM, Pratchett M, Schoepf V, Torda G, Wilson SK. 2018. Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* 359: 80-83.
27. Intergovernmental Panel on Climate Change. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Geneva, Switzerland: Intergovernmental Panel on Climate Change; 2014.
28. Jacobson MZ. Clean grids with current technology. 2016. *Nature Climate Change* 6: 441. 25 January 25, 2016.

29. Johnston F, Hanigan I, Henderson S, Morgan G, Bowman D. 2011. Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994-2007. *Environmental Research* 111: 811-816. Doi: 10.1016/j.envres.2011.05.007
30. Karl TR, Melillo JM, Peterson TC, eds. *Global climate change impacts in the United States*. New York: Cambridge University Press: 2009.
31. Karl TR, Arguez A, Huang B, Lawrimore JH, McMahon JR, Menne MJ, Peterson TC, Vose RS, Zhang HM. 2015. Possible artifacts of data biases in the recent global surface warming hiatus. *Science* 348: 1469-1472. Doi: 10.1126/science.aaa5632
32. Kopp RE, Horton RM, Little CM, Mitrovica JX, Oppenheimer M, Rasmussen DJ, Strauss BH, Tebaldi C. 2014. Probabilistic 21st and 22nd century sea-level projections at a global network of tide-gauge sites. *Earth's Future* 2: 383-40618.
33. Lelieveld J, Evans JS, Fnais M, Giannadaki D, Pozzer A. 2015. The contribution of outdoor air pollution sources to premature mortality on a global scale. *Nature* 525: 367-371. Doi: 10.1038/nature15371
34. Levinson DH and Fettig C. 2014. Climate change: Overview of data sources, observed and predicted temperature changes, and impacts on public and environmental health. In *Global Climate Change and Public Health*, Pinkerton KE and Rom WN, eds, New York: Humana Press, 2014.
35. Liu-Helmersson J, Stenlund H, Wilder-Smith A, Rocklov J. 2014. Vectorial capacity of *Aedes Aegypti*: Effects of temperature and implications for global dengue epidemic potential. *PLOS ONE* 9(3): e89783. Doi: 10.1371/journal.pone.0089783
36. Luthi D, LE Floch M, Bereiter B, Blunier T, Barnola JM, Siegenthaler U, Raynaud D, Jouzel J, Fischer H, Kawamura K, Stocker TF. 2008. High-resolution carbon dioxide concentrations record 650,000-800,000 years before present. *Nature* 453: 379-382. Doi:10.1038/nature06949
37. MacDonald AE, Clack CTM, Alexander A, Dunbar A, Wilczak J, and Xie Y. 2016. Future cost-competitive electricity systems and their impact on US CO₂ emissions. *Nature Climate Change* 6: 526-531. DOI: 10.1038/NCLIMATE2921
38. Markandya A, Armstrong BG, Hales S, Chiabai A, Criqui P, Minna S, Tonne C, Wilkinson P. 2009. Public health benefits of strategies to reduce greenhouse-gas emissions: low-carbon electricity generation. *Lancet* 374: 2006-2015.
39. Mazdiyasi O, AghaKouchak A, Davis SJ, Madadgar S, Mehran A, Ragno E, Sadegh M, Sengupta A, Ghosh S, Dhanya CR, Niknejad M. 2017. Increasing probability of mortality during Indian heat waves. *Sci Adv* 3: e1700066. Doi: 10.1126/sciadv.1700066.
40. McIver L, Kim R, Woodward A, Hales S, Spickett J, Katscherian D, Hashizume M, Honda Y, Kim H, Iddings S, Naicker J, Bambrick H, McMichael AJ, Ebi KL. 2016. Health impacts of climate change in Pacific Island countries: A regional assessment of vulnerabilities and adaptation priorities. *Environ Health Perspect* 124: 1707-1714.
41. McKibben B. 2012. Global warming's terrifying new chemistry. *The Rolling Stone*, July 19, 2012.
42. McMichael AJ. 2013. Globalization, Climate Change, and Human Health. *N Engl J Med* 368: 1335-43. DOI: 10.1056/NEJMr1109341.
43. Millar RJ, Fuglesteit JS, Friedlingstein P, Rogelj J, Grubb MJ, Matthews HD, Skeie RB, Forster PM, Frame DJ, Allen MR. 2017. Emission budgets and pathways consistent with limiting warming to 1.5°. *Nature Geoscience* 10: 741-747. doi: 10.1038/NGE03031.
44. Molina M, Zaelke D, Sarma KM, Andersen SO, Ramanathan V, Kaniaru D. 2009. Reducing abrupt climate change risk using the Montreal Protocol and other regulatory actions to complement cuts in CO₂ emissions. *Proc Natl Acad Sci USA* 106: 201616-21.
45. Montzka SA, Dlugokencky EJ, Butler JH. 2011. Non-CO₂ greenhouse gases and climate change. *Nature* 476: 43-50. Doi: 10.1038/nature10322
46. Mora C, Dousset B, Caldwell IR, Powell FE, Geronimo RC, Bielecki CR, Counsell CWW, Dietrich BS, Johnston ET, Louis LV, Lucas MP, McKenzie MM, Shea AG, Tseng H, Giambelluca TW, Leon LR, Hawkins E, Trauernicht C. 2017. Global risk of deadly heat. *Nature Climate Change* 7: 501-506. DOI: 10.1038/NCLIMATE3322.
47. Morabia A, Benjamin GC. 2017. Preparing and rebuilding after natural disasters: A new public health normal! *American Journal of Public Health*. 108: 9-10. Doi: 10.2105/AJPH.2017.304202.
48. Morbidity and Mortality Report. Deaths Associated with Hurricane Sandy October-November 2012. *May 24, 2013/ 62(20); 393-397.*

49. Moritz C and Agudo R. 2013. The future of species under climate change: Resilience or decline? *Science* 341: 504-508. Doi: 10.1126/science.1237190
50. National Academies of Science, Engineering and Medicine. 2016. Attribution of extreme weather events in the context of climate change. Washington, DC: The National Academies Press. DOI: 10.17226/21852.
51. O'Neill MS, Zanobetti A, Schwartz J. 2005. Disparities by race in heat-related mortality in four US cities: The role of air conditioning prevalence. *J Urban Health* 82: 191-197.
52. Pacala S and Socolow R. 2004. Stabilization wedges: Solving the climate problem for the next 50 years with current technologies. *Science* 305: 968-972.
53. Park C-E, Jeong S-J, Joshi M, Osborn TJ, Ho C-H, Piao S, Chen D, Liu J, Yang H, Park H, Kim B-M, Feng S. 2018. Keeping global warming within 1.5° C constrains emergence of aridification. *Nature Climate Change* 8: 70-74. Doi.org/10.1038/s41558-017-0034-4
54. Patz J. 2014. Climate variability and change: Food, water, and societal impacts. In *Global Climate Change and Public Health*, Pinkerton KE and Rom WN, eds, New York: Humana Press, 2014.a
55. Patz JA, Frumkin H, Holloway T, Vimont DJ, Haines A. 2014. Climate Change. Challenges and Opportunities for Global Health. *JAMA* 312(15): 1565-1580. Doi: 10.1001/jama.2014.13186.b
56. Pershing AJ, Alexander MA, Hernandez CM, Kerr LA, Le Bris A, Mills KE, Nye JA, Record NR, Scannell HA, Scott JD, Sherwood GD, Thomas AC. 2015. Slow adaptation in the face of rapid warming leads to collapse of the Gulf of Maine cod fishery. *Science* 350: 809-812. Doi: 10.1126/science.aac9819
57. Rice MB, Thurston GD, Balmes JR, Pinkerton KE. 2014. Climate Change. A global threat to cardiopulmonary health. *Am J Respir Crit Care Med* 189: 512-519.
58. Rom WN, Pinkerton KE, Martin WJ, Forastriere F. 2008. Global warming: A challenge to all American Thoracic Society members. *Am J Respir Crit Care Med*. 177: 1053-7.
59. Rom WN, Evans L, and Uppal A. 2013. The sentinel event of climate change. Hurricane Sandy and its consequences for pulmonary and critical care medicine. *Am J Respir Crit Care Med* 187: iii-iv.
60. Rom WN and Pinkerton KE. 2014. Introduction: Consequences of Global Warming to the Public's Health. In Pinkerton KE and Rom WN eds., *Global Climate Change and Public Health*. New York: Humana Press, 2014.
61. Solomon S, Plattner G-K, Knutti R, Friedlingstein P. 2009. Irreversible climate change due to carbon dioxide emissions. *Proc Natl Acad Sci USA* 106: 1704-1709. Doi: 10.1073/pnas.0812721106
62. Stott P. 2016. How climate change affects extreme weather events. *Science* 352: 1517-1518.
63. Taylor CM, Belusic D, Guichard F, Parker DJ, Vischel T, Bock O, Harris PP, Janicot S, Klein C, Panthou G. 2017. Frequency of extreme Sahelian storms tripled since 1982 in satellite observations. *Nature* 544: 475-478. doi:10.1038/nature22069.
64. The White House. November 2016. United States Mid-Century Strategy for deep decarbonization.
65. US Global Change Research Program. 2014. Climate change impacts in the United States. Washington, DC: US Global Change Research Program; 2014. Available from: <http://nca2014.globalchange.gov/>
66. Watts N, Adger WN, Agnolucci P, et al. 2015. Health and climate change: policy responses to protect public health. *Lancet* 24 Jun 2015; DOI: 10.1016/S0140-6736(15)60854-6.
67. Watts N, Amann M, Ayeb-Karlsson S, et al. 2017. The *Lancet* Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet* 1-50. doi.org/10.1016/S0140-6736(17)32464-9.
68. West JJ, Smith SJ, Silva RA, Naik V, Zhang Y, Adelman Z, Fry MM, Anenberg S, Horowitz LW, Lamarque J-F. 2013. Co-benefits of mitigating global greenhouse gas emissions for future air quality and human health. *Nature Climate Change* 3(10): 885-889. Doi 10.1038/NCLIMATE2009.
69. Whitmee S, Haines A, Beyrer C, et al. 2015. Safeguarding human health in the Anthropocene epoch: report of the Rockefeller Foundation-Lancet Commission on planetary health. *Lancet* 1-55. doi.org/10.1016/S0140-6736(15)60901-1.
70. Zeebe RE, Ridgwell A, Zachos JC. 2016. Anthropogenic carbon release rate unprecedented during the past 66 million years. *Nature Geoscience* 9:325-329. DOI: 10.1038/NGEO2681.